



# **Metropolitan Water Reclamation District of Greater Chicago**

**Welcome to the June  
Edition of the 2021  
M&R Seminar Series**


## NOTES FOR SEMINAR ATTENDEES

- All attendees' audio lines have been muted to minimize background noise.
- A question and answer session will follow the presentation.
- Please use the "Chat" feature to ask a question via text to "All Panelists".
- The presentation slides will be posted on the MWRD website after the seminar.
- This seminar has been approved from the ISPE for one PDH, and pending approval from the IEPA for one TCH. Certificates will only be issued to participants who attend the entire presentation.

## **Wei Zheng, Ph.D.**

Principal Research Scientist


Illinois Sustainable Technology Center/Prairie Research Institute  
University of Illinois at Urbana-Champaign



Dr. Wei Zheng is a principal research scientist at the Illinois Sustainable Technology Center (ISTC)/Prairie Research Institute (PRI) at the University of Illinois at Urbana-Champaign (UIUC). He is also an adjunct faculty member at UIUC. He joined the ISTC/PRI/UIUC in 2008. Prior to joining the ISTC, he worked as a post-doc and soil scientist at USDA-ARS, Salinity Laboratory and University of California, Riverside. He earned a Ph.D. in Environmental Chemistry from Zhejiang University (China). He has expertise in environmental science and environmental chemistry with extensive experience on the fate and transport of pesticides and emerging contaminants including pharmaceuticals and personal care products (PPCPs) and per- and polyfluoroalkyl substances (PFAS), remediation techniques for chemical pollutants, and production and utilization of biochar. Currently, he has multiple on-going projects funded from USDA, USEPA, DOE and IL NREC, which are related to biochar.



# *Biochar in Soil and Environmental Applications*



**Wei Zheng**

*Illinois Sustainable Technology Center  
University of Illinois at Urbana-Champaign  
[weizheng@illinois.edu](mailto:weizheng@illinois.edu)*



## Illinois Sustainable Technology Center (ISTC)

The Illinois Sustainable Technology Center (ISTC) is a Division of the Prairie Research Institute (PRI) at the University of Illinois (U of I). ISTC (formerly Illinois Waste Management Research Center) has been providing **research, scientific expertise, and data that benefit the environment, economy, and people of Illinois and beyond**. It used to be Illinois State Agency.

Current ISTC missions include:

- Research
- Technical assistance
- Technique demonstration
- Communication

<http://www.istc.illinois.edu/>





# Acknowledgments



- Sarmila Katuwal
- Nathan Pflugmacher
- Marisa Moreno
- Mahelet Maru
- Siming Yang
- John Scott
- Wiles, K.
- Murali, D.
- Manandhar, J
- Joe Pickowitz
- Dr. Sharma, B.K.
- Dr. Richard Cooke
- Dr. Rajagopanlan, K.

## MWRD

- Dr. Olawale Oladeji
- Dr. Guanlong Tian
- Dr. Albert Cox
- Jacob Baylor
- Dr. Edward Podczerwinski
- Dr. Dominic Brose
- Lauren Lurkins
- Elaine Stone
- Dr. Lakhwinder Hundal



# What is Biochar?

- **Biochar** is a carbon-enriched material produced from **biomass** under a limited supply of oxygen. “Not a high-tech product”.



**Biomass**




**Pyrolysis**

**Syngas**  
 $H_2$ ,  $CO$ ,  $CO_2$



**Bio-oil**



**Biochar**



# Biochar vs Other Carbons

- Biochar is similar in its appearance to charcoal, char, activated carbon, and other black carbon materials. While biochar is produced from **waste biomass**, it can provide the following benefits.
  - ❖ *Waste reduction*
  - ❖ *Energy production*
  - ❖ *Contaminant sorbent*
- The difference of biochar and carbon-rich materials:  
Biochar is produced from biomass with the intent to deliberately apply to soil for:
  - ❖ *Soil Amendment*
  - ❖ *Carbon Sequestration*





# Biochar Applications in Soil and Environment

Overall, biochar applications in soil and environment can provide multiple valuable benefits, such as E<sup>3</sup> (Energy, Environmental, Economic) profits.

Our biochar studies focus on:


- **Soil amendment** ←
- **Carbon sequestration**
- **Environmental Applications**
  - ❖ **Nutrient capture**
  - ❖ **Emerging contaminant removal**



## Biochar as a Soil Amendment

Once biochars are added to soil, they can **boost soil fertility and improve soil quality** by

- ❖ trapping moisture
- ❖ helping the soil hold nutrients
- ❖ attracting more beneficial fungi and microbes
- ❖ improving cation exchange capacity (CEC)
- ❖ raising soil pH
- ❖ others....





## Biochar & Slash-and-Burn

- **Slash-and-burn** is a specific functional element of certain farming practices, often shifting cultivation systems. Before artificial fertilizers were available, it was traditional fertilization method. Burning removes the vegetation and may release a pulse of nutrients to fertilize the soil.
- **Slash-and-char** is an alternative to slash-and-burn that has a lesser effect on the environment. Switching from “slash-and-burn” to “slash-and-char” can prevent the rapid deforestation and subsequent degradation of soils. It results in the creation of biochar, which can then be mixed with biomass such as crop residues, food waste, manure or other, and buried in the soil to bring about the formation of terra preta. Terra preta is one of the richest soils n on the planet.






## Amazonian Dark Soil---*terra preta*

**Black  
topsoil**

the charcoal  
deposit site

**Yellow  
underlying  
material**



*fertile*



*nutrient-poor*

Comparison of profiles of *terra preta* and adjacent soils

----From: IBI website

## Biochar as a Soil Amendment

- However, biochar additions to soils have not consistently resulted in increased crop yields. It has been reported that approximately 50% of the studies showed short-term yield increases after biochar soil application, 30% observed no significant differences, and 20% noted negative yield or growth impact.
- Another barrier for biochar large scale application is cost --- improving biochar values to cover the cost.



**No biochar**

**2% biochar**

**4% biochar**

**8% biochar**



# Projects on **Biochar & Chemical Fertilizer Reduction**

This study is to investigate if the use of biochar as a soil amendment in **Illinois rich soils** can reduce chemical fertilizer use while at the same time maintaining or increasing crop yields. To achieve this goal, three tasks were undertaken in the project:

➤ **Biochar production and characterization:**

Biochar production through a low-temperature slow pyrolysis technique from a variety of waste biomass.

➤ **Sorption of nutrients by biochar:**

The sorption kinetics and mechanisms of  $\text{NH}_4^+$  and  $\text{PO}_4^{3-}$  removal by biochar were investigated.

➤ **Greenhouse and Field trial:**

Demonstrate the efficacy of biochar as a simple soil amendment as measured by crop yields and lowered fertilizer use in Illinois rich soils.

*Illinois Hazardous Waste Funds* & *Russell and Helen Dilworth Memorial Fund*





## Biochar Application in Greenhouse

We used a radish as a target plant to evaluate the effect of different application rates of biochars and nitrogen fertilizers on crop yields.

- Biochar application rates: 0, 2%, 4%, 8% (w/w).
- Nitrogen fertilize ( $\text{NH}_4\text{NO}_3$ ) rates: 0, 25, 50, and 100 lbs N/acre.





## Biochar Application in Greenhouse

Only biochar



No biochar

2% biochar

4% biochar

8% biochar


In the absence of fertilizer, biochar use resulted in no improvement in either radish plant growth or crop yields.

# Biochar Application in Greenhouse

N fertilizer application rates



Biochar application rates





# Biochar Application in Illinois Corn Fields







April



May-June



June-July




Aug.- Oct.







# Biochar Application in Corn Field



Nitrogen Fertilizer	0	50 %	100 %
No Biochar	139.3 <sup>a</sup>	174.3 <sup>a</sup>	173.0 <sup>a</sup>
Biochar-A	164.6 <sup>b</sup>	213.7 <sup>b</sup>	239.8 <sup>b</sup>
Biochar-B	170.9 <sup>b</sup>	194.2 <sup>b</sup>	201.3 <sup>a</sup>

## Sorption of $\text{NH}_4^+$ on Various ISTC's Biochars

Ammonium ion  $C_s$  (mmol/g)





# Biochar Applications in Soil and Environment

Overall, biochar applications in soil and environment can provide multiple valuable benefits, such as E<sup>3</sup> (Energy, Environmental, Economic) profits.

Our biochar studies focus on:

- **Soil amendment**
- **Carbon sequestration**
- **Environmental Applications**
  - ❖ **Nutrient capture**
  - ❖ **Emerging contaminant removal**



# Global Climate Change and Global Warming

**Positive proof of global warming.**



**18th Century 1900 1950 1970 1980 1990**


**Global Climate change is a big challenge**

Large emissions of greenhouse gases, primarily from CO<sub>2</sub>




**Global atmospheric concentration of CO<sub>2</sub> (since industrial revolution)**

Global atmospheric concentration of CO<sub>2</sub>







# Climate mitigation from “nature-based” activities





# How does Biochar Sequester Atmospheric Carbon?

Biochar is one of feasible solutions to reduce the growing levels of CO<sub>2</sub> in the atmosphere.

The natural carbon cycle is carbon neutral. If we use pyrolysis to produce biochar from waste biomass, we can lock up 50% biomass carbon into biochar. Considering carbon in biochar can store in soils for a long term, the net process is carbon negative. Therefore, biochar production from biomass and its application into soils has a great potential to alleviate climate change.





**A****B**

## Carbon Footprint Analysis

### Schematics for biomass or bio-char remaining after charring and decomposition in soil.

(A) C remaining from biomass decomposition after 100 years from IPCC; C remaining after charring or pyrolysis; bio-char C remaining after decomposition.  
 (B) range of biomass C remaining after decomposition of crop residues and estimation of bio-char.

**From:** Lehmann et al. 2006  
 in *Mitigation and Adaptation Strategies for Global Change*



## Projects on **Improving Biochar Carbon Sequestration Capacity**

- The **technique** is to pre-treat biomass by mixing some passivator-like materials to retain more biomass carbon in biochar during pyrolysis.
- The **objective** of this research project is to use chemical modification methods to strengthen carbon retention and stability in biochar.

This study is being supported by  
**Russell and Helen Dilworth Memorial Fund**



# Effect of Chemical Additives on Carbon Retention




**Six P-rich chemicals or passivator-like materials** were chosen as chemical additives in this study: triple superphosphate (TSP), phosphate rock tailing (PRT), bonemeal (BM), clay,  $\text{H}_3\text{PO}_4$ , and  $\text{H}_3\text{BO}_3$ .

## Triple Superphosphate (TSP)


**Triple superphosphate (TSP)** is P-containing fertilizer. Also, it is an environmentally friendly and low-cost chemical additive. Thus, TSP was chosen as a preferred additive for the following studies.

TSP is produced from “phosphate rock”.






# Effect of TSP Addition Amounts on Carbon Retention



switchgrass




# Interaction Mechanism of TSP and Biomass-Carbon During Pyrolysis



A new peak at  $2\theta^\circ=26.6$  which corresponds to the C-P chemical bond was observed in all P-chemical modified biochars



# Interaction Mechanism of TSP and Biomass-Carbon During Pyrolysis



X-ray photoelectron spectroscopy (XPS)

The appearance of the peak at ~135.0 eV in TSP modified biochar is attributed to the formation of the  $C-O-PO_3$  chemical bond.

XPS spectra of P 2p



# Biochar Stability

## Schematics for biomass or bio-char remaining after charring and decomposition in soil.

(A) C remaining from biomass decomposition after 100 years from IPCC; C remaining after charring or pyrolysis; bio-char C remaining after decomposition.


(B) range of biomass C remaining after decomposition of crop residues and estimation of bio-char.

**From:** Lehmann et al. 2006  
in *Mitigation and Adaptation Strategies for Global Change*






# Biochar Thermochemical Stability by TGA



Compared to biochars produced from biomass only, TSP-modified biochars need much higher temperatures for decomposition based on thermogravimetric analysis (TGA), suggesting the TSP- addition could increase biochar thermochemical stability.


# Biological Mineralization of TSP-Modified Biochar in Soil



CO<sub>2</sub>-C cumulative emissions from soils amended with biomass, biochar, and TSP-modified biochar produced from switchgrass as well as soil alone (as a control study)



# Carbon Footprint Analysis of Biochar




## P Leaching from TSP-Modified Biochars


<b>Biomass</b>	<b>Biochar</b>	<b>Leached P (mg) per g solid</b>
<b>Sawdust biochars</b>	$BC_{\text{saw}}$	0.153
	$BC_{\text{saw}}\text{-TSP}$	7.07
<b>Switchgrass biochars</b>	$BC_{\text{swi}}$	0.110
	$BC_{\text{swi}}\text{-TSP}$	6.18
<b>TSP</b>		101



# TSP-Modified Biochar for Slow-Release Fertilizer ??



# TSP-Modified Biochar for Greenhouse Trial



**Application rate = 0.25% (significantly different,  $p=0.01$ )**



# Biochar Applications in Soil and Environment

Overall, biochar applications in soil and environment can provide multiple valuable benefits, such as E<sup>3</sup> (Energy, Environmental, Economic) profits.

Our biochar studies focus on:

- **Soil amendment**
- **Carbon sequestration**
- **Environmental Applications**
- ❖ **Nutrient capture**
- ❖ **Emerging contaminant removal**



## Nutrient Loss from Tile Drainage


- In the Midwestern U.S., subsurface tile drainage systems have been widely used in agricultural fields to remove excess water from the soil profile to increase crop production and promote soil conservation.
- However, tile drainage can deliver large quantities of nutrients, primarily nitrate-nitrogen ( $\text{NO}_3\text{-N}$ ) and phosphorus, from agricultural fields into receiving watersheds. It is of particular concern in the Midwest since this nutrient delivery has been identified as a major contributor to the growing hypoxia zone in the Gulf of Mexico. Excess amounts of phosphorus in marine and freshwater systems is often identified as a primary perpetrator for the harmful algal blooms.





# Peak Phosphorus

- Recently, phosphorus depletion or “Peak Phosphorus” has been gaining increased public attention since the world is running short of phosphorus ore for chemical fertilizers.
- It is critical to develop best management practices (BMPs) that can reduce phosphorus losses from fields and recover lost phosphorus for agricultural application.



## Research Idea

- The idea is to produce designer biochars to simultaneously capture ammonium nitrogen ( $\text{NH}_4^+\text{-N}$ ) and phosphate ions from tile drainage water.
- The nutrient-captured biochars will be used as a slow-release fertilizer to effectively keep nutrients in agricultural loop, minimize their losses, and enhance soil fertility.





## Removal Mechanisms of Phosphate & Ammonium by Biochar



Analyte Units	Biochar after DI H <sub>2</sub> O washing	Biochar no washing
	mg/L	mg/L
<b>Sodium*</b>	< 0.6	< 0.6
<b>Potassium*</b>	7.7	15
<b>Calcium*</b>	2.8	3.8
<b>Beryllium</b>	< 0.002	< 0.002
<b>Boron</b>	0.089	0.094
<b>Magnesium</b>	1.5	2.3
<b>Aluminum</b>	0.075	0.084
<b>Silicon</b>	2.0	1.0
<b>Titanium</b>	0.0027	0.0022
<b>Vanadium</b>	< 0.001	< 0.001
<b>Chromium</b>	0.0032	0.0033
<b>Manganese</b>	0.0077	0.0055
<b>Iron*</b>	< 0.1	0.11
<b>Zeta potential (ζ)</b>	<b>-21.9±3.8</b>	<b>-22.3±6.9</b>

➤ **Electrostatic adsorption**  
Biochar with negatively charged surface



# Designer Biochar Production

- The surfaces of most biochars are negative charge, so they are generally ineffective to remove phosphorus, especially anionic phosphate ions ( $\text{PO}_4^{3-}$ ,  $\text{HPO}_4^{2-}$  and  $\text{H}_2\text{PO}_4^-$ ).
- Designer biochars are produced by pyrolysis of biomass pre-treated with calcium-enriched materials lime sludge (main component  $\text{CaCO}_3$  with high Mg contents) and gypsum (main component  $\text{CaSO}_4$ ). They are being used as agricultural lime. Lime sludge is a by-product from the softening process in water treatment plants and gypsum is a by-product from power plants.






- The **objectives** of these research projects are to generate designer biochars to effectively capture nutrients (especially for dissolved phosphorus) from subsurface tile drainage, recycle nutrient-captured biochars as a slow released fertilizer, and keep phosphorus in the agricultural loop.

### Tasks:

- Create designer biochars by pyrolysis of biomass pre-treated with calcium-enriched materials (e.g., lime sludge and gypsum)
- Conduct a laboratory experiment to optimize the biochar production conditions by evaluating their sorption capacities for ammonium and phosphate ions.
- Manufacture designer biochars to capture nutrients from subsurface tile drainage by constructing biochar-sorption-channel in fields.
- Recycle nutrient-captured biochars and apply them as a slow-release fertilizer to improve soil quality and crop yields
- Develop and scale-up a

# Ammonium Removal by Designer Biochars






S/L=1:100 in 0.5 mm NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> solution


**Ammonium sorption capacities of designer biochars produced from different temperatures**



# Optimize Designer Biochar Production Conditions to Improve Phosphorus Sorption




# Phosphorus Removal by Different Adsorbents




Comparison of phosphate removal by different adsorbents

S/L=1:250 in 50mg/L phosphate solution



# Adsorption Mechanisms — XRD Spectrum








XRD patterns of the unmodified biochar, the designer biochar, and the P-laden designer biochar



# Adsorption Mechanisms of Phosphate by Designer Biochar

Adsorption of P could be controlled by multiple mechanisms

- *Intrapartical diffusion on designer biochar (physical adsorption)*
- *Surface deposition reactions on designer biochar:*



- *Activate precipitation of P by designer biochar (i.e., high pH)*

Alkalinity is one of the most influential properties for most biochars because their surfaces contain a large amount of alkaline functional groups



# Production of Designer Biochar Pellets



+



**Pelletizing**




**Pyrolysis**



**Designer  
Biochar**



# Bioreactor and P-Sorbing Filter




The field test is being conducted at Fulton County, IL. Currently, three 11.5-acre drainage systems have been installed at the MWRD research site at Fulton County. Each drainage system includes three bioreactor compartments with controlled structure and space to hold P-sorbing filters



# Designer Biochar to Capture Nutrients from Tile Drainage in Fields

**Three treatments** are being implemented in triplicate in the existing drainage systems:

- (1) The designer biochars was packed into the P-sorbing filters
- (2) A commercial polymer-based phosphorus sorbent-RecoNP, which was provided by one of collaborators (Dr. Hundal), was packed into the P-sorbing filters; and
- (3) A slag material was packed into the P-sorbing filters.




# Designer Biochar to Capture Nutrients from Tile Drainage in Fields



Polymer Sorbent


Iron Material

Design Biochar





## Nutrient-captured biochar application





# Biochar-Sorption-Chamber Installation





# Nutrient-Laden Biochar for Slow Release Fertilizer



**ISTC Slow-Release Fertilizer**

# Biochar Applications in Soil and Environment

Overall, biochar applications in soil and environment can provide multiple valuable benefits, such as E<sup>3</sup> (Energy, Environmental, Economic) profits.

Our biochar studies focus on:


- **Soil amendment**
- **Carbon sequestration**
- **Environmental Applications**
  - ❖ **Nutrient capture**
  - ❖ **Emerging contaminant removal**






# Removal of Pharmaceuticals and Personal Care Products by Biochar

- Currently, pharmaceuticals and personal care products termed PPCPs are receiving great public attentions as emerging contaminants.
- PPCPs have been frequently detected in the lake, river, and stream in the US because they are widely used in human and veterinary clinical practices.
- Because biochar has a high adsorption capacity, it has potential to remove these emerging contaminants




# Sorption of Florfenicol by Biochars Produced from Different Waste Biomass






# Sorption of Steroid Hormones by Biochars Produced from Wood Waste



# Per- and Polyfluoroalkyl Substance (PFAS)

- Aqueous Film Forming Foams
- Stain-Water Resistant Textiles
- Plastics
- Personal Care Products
- Metal Coatings
- Paints
- Cleaning Products
- Oil and Mining Additives
- Food Packaging





# PFAS


**PFAS are “Forever Chemicals”.** They build up in our bodies and never break down in the environment. Currently, they widely occur in the environment. For example, up to 110 million people may be drinking PFAS-containing water.

**PFAS are highly bioaccumulated chemicals.** Today nearly all Americans, including newborn babies, have PFAS in their blood. A growing body of science has found that there are potential adverse health impacts associated with PFAS exposure, including liver damage, thyroid disease, decreased fertility, high cholesterol, obesity, hormone suppression and cancer.



## Per- and Polyfluoroalkyl Substance (PFAS)

The occurrence of per- and polyfluoroalkyl substances (PFASs) in the environment, coupled with their known adverse effects on public health, has been recognized as a widespread issue. As a result of legacy use of aqueous film forming foam (AFFF) in firefighting activities, the U.S. Department of Defense (DoD) may have hundreds of sites contaminated with PFASs. It is critical to develop cost-effective and efficient strategies to manage and remediate these PFAS contaminated sites.



In May 2016, the U.S. Environmental Protection Agency (EPA) promulgated drinking water Health Advisory Levels (HALs) for PFOS and PFOA (2, 16). The current HAL for drinking water is 70 ng/L for the sum of PFOS and PFOA, or either compound individually



# PFAS Contamination Sources and their Fate/Transport in the Environment



# PFAS Fate and Transport Proposals




- Funding Opportunity Title and Number: Practical Methods to Analyze and Treat Emerging Contaminants (PFAS) in Solid Waste, Landfills, soils, and Groundwater to Protect Human Health and protect Human Health and the Environment, EPA-G2018-STAR-B1

- Project Title: **Linkage between physiochemical properties of biosolids-associated PFASs and their behavior in soil-plant systems**




Funding Opportunity Title and Number: National Priorities: Research on PFAS Impacts in Rural Communities and Agricultural Operations, EPA-G2020-ORD-B1

Project Title: **Systematic Understanding of PFAS Sources and Mobility to Water Resources from Biosolids Land Application and Water Reuse in the US Rural Areas**



# NIEHS P42 Superfund Center

The National Institute of Environmental Health Science (NIEHS) proposes the continuation of the Superfund Hazardous Substance Research and Training Program (P42) to address the broad, complex health and environmental issues that arise from hazardous waste sites.




Outline of UIUC's NIEHS Superfund Five Research Cores




# PFAS Remediation/Treatment Techniques

## Generate a novel biochar to remove PFAS from contaminated water

- *Biochar* is a carbon-enriched material produced from biomass under a limited supply of oxygen. “Not a high-tech product”.
- Biochar is similar in its appearance to charcoal, char, activated carbon, and other black carbon materials.
- The difference of biochar and carbon-rich materials:  
Biochar is produced with the intent to deliberately apply to soil to:
  - ❖ *Sequester carbon*
  - ❖ *Improve soil quality*



# PFAS Remediation/Treatment Techniques



Comparison of PFAS sorption capacities by biochar, designer biochar, and activated carbon







## Remediation of PFAS Contaminated Biosolids and AFFF-Impacted Soils

# PFAS Remediation/Treatment Techniques

Develop innovation mitigation techniques to prevent PFAS leaching from contaminated biosolids and AFFF-impacted soils:

- **Biosolid-based biochar**: Use of low temperature (<450 °C) pyrolysis to entrap PFAS during biosolid carbonization, stabilize them in biosolid-based biochar, and thereby reduce their availability to leaching and plant uptake.
- Generate an innovative designer biochar and compost it with contaminated biosolids and/or AFFF-impacted soils to constrain the mobility of PFAS and mitigate their leaching




Figure 9. PFOA leaching from biosolids and Biosolids-based biochar




# PFAS Remediation/Treatment Techniques

Develop innovation mitigation techniques to prevent PFAS leaching from contaminated biosolids and AFFF-impacted soils:


- Biosolid-based biochar: Use of low temperature (>450 °C) pyrolysis to entrap PFAS during biosolid carbonization, stabilize them in biosolid-based biochar, and thereby reduce their availability to leaching and plant uptake.
- Generate an innovative ***designer biochar*** and compost it with contaminated biosolids and/or AFFF-impacted soils to constrain the mobility of PFAS and mitigate their leaching

PFA removal efficiency of different biochar





# Summary



# What is Biochar?

- **Biochar** is a carbon-enriched material produced from **biomass** under a limited supply of oxygen. “Not a high-tech product”.



**Biomass**

**Waste biomass** includes:

- Forest and agricultural residues.
- Yard wastes



**Pyrolysis**

**Syngas**  
 $H_2$ ,  $CO$ ,  $CO_2$



**Bio-oil**



**Biochar**





# Biochar as a Soil Amendment

Once biochars are added to soil, they can improve soil quality.

When used as a soil amendment, biochar can boost soil fertility and improve soil quality by


- ❖ trapping moisture
- ❖ helping the soil hold nutrients
- ❖ attracting more beneficial fungi and microbes
- ❖ improving cation exchange capacity (CEC)
- ❖ raising soil pH
- ❖ others....






# Carbon Sequestration vs Global Warming

**Positive proof of global warming.**




18th  
Century      1900      1950      1970

**Climate change**  
Large emissions of  
greenhouse gases, primarily  
from CO<sub>2</sub>





**Global atmospheric concentration  
of CO<sub>2</sub> (since industrial revolution)**



# Bioreactor and Biochar-Sorption-Channel

## Nutrient-captured biochar application







# PFAS Remediation/Treatment Techniques

## Destruction of PFAS during Designer Biochar Hydrothermal Regeneration

The PFAS-captured biochar will be reactivated using hydrothermal treatment to recover the sorbent under mild regeneration temperatures while at the same time to defluorinate and decompose the PFAS.







# PFAS Remediation/Treatment Techniques

Develop innovation mitigation techniques to prevent PFAS leaching from contaminated biosolids and AFFF-impacted soils:

- **Biosolid-based biochar**: Use of low temperature (>450 °C) pyrolysis to entrap PFAS during biosolid carbonization, stabilize them in biosolid-based biochar, and thereby reduce their availability to leaching and plant uptake.
- Generate an innovative **designer biochar** and compost it with contaminated biosolids and/or AFFF-impacted soils to constrain the mobility of PFAS and mitigate their leaching



# Questions??



*Wei Zheng*  
[weizheng@Illinois.edu](mailto:weizheng@Illinois.edu)  
217-333-7276